

Tri-D

[Cholecalciferol]

INJECTION

COMPOSITION

Each 1ml ampoule contains:
Cholecalciferol (Vit. D₃) BP.....5mg (200,000 IU)

DOSAGE FORM

Drinkable and injectable solution in 1ml ampoule.

ROUTE OF ADMINISTRATION

I.M route. The ampoule can also be administered by oral route.

PHARMACOTHERAPEUTIC GROUP

Vitamin D. (Fat Soluble Vitamin.)

INDICATIONS

This medicinal product contains Vitamin D, which is involved in bone fixation of calcium. It is indicated in prevention and treatment of Vitamin D deficiencies.

WARNING!

This drug must not be used in the following cases:

- Hypersensitivity to any of the ingredients, mainly to Vitamin D
- Hypercalcemia (abnormally high blood calcium levels)
- Hypercalciuria (excessive urinary elimination of calcium)
- Calcium lithiasis (kidney stones)

IF YOU ARE IN DOUBT, YOU MUST ASK YOUR PHYSICIAN.

SPECIAL WARNINGS

In order to avoid any Vitamin D over dosage, you should ask your doctor's advice in case of concomitant treatment with drugs containing this vitamin.

If high or repeated doses of Vitamin D are administered or if high doses of calcium are associated, it is necessary to monitor calcium levels in blood and urine.

PRECAUTIONS FOR USE

If you are in doubt do not hesitate to ask your physician.

INTERACTIONS

In order to avoid possible interactions between several drugs you must always tell your physician about any other treatments you are taking.

PREGNANCY AND BREAST-FEEDING

This medicinal product can be prescribed during pregnancy or lactation if necessary. However it is preferable to consult your doctor or pharmacist before using drugs.

DOSAGE

In all cases follow your physician's prescription.

USUAL DOSAGES ARE:

INFANTS

- **INFANTS RECEIVING VITAMIN ENRICHED MILK:** 1/2 ampoule (i.e. 100,000 IU) every 6 months.

- **NURSED INFANTS OR INFANTS NOT RECEIVING VITAMIN D ENRICHED MILK OR YOUNG CHILDREN UP TO 5 YEARS OF AGE:** 1 ampoule (200,000 IU) every 6 months.

ADOLESCENTS

1 ampoule (200,000 IU) every 6 months during winter.

ELDERLY

1/2 ampoule (i.e. 100,000 IU) every 3 months.

PREGNANCY

1/2 ampoule (i.e. 100,000 IU) from 6th or 7th month of pregnancy.

Digestive disorders, concomitant treatment with anti-epileptics, other particular conditions not described above: 1/2 or 1 ampoule every 3 or 6 months.

VITAMIN-D DEFICIENCY

1 ampoule (i.e. 200,000 IU), which can be renewed once 1 to 6 months later.

THIS MEDICINE HAS BEEN PRESCRIBED TO YOU PERSONALLY FOR A PARTICULAR CONDITION:

- It can not be adapted for another condition
- Do not recommend it to another person

OVERDOSE

The following symptoms may occur: headache, fatigue, slimming, growth retardation, nausea, vomiting, excess of urine, intense thirst, arterial hypertension.

INFORM YOUR PHYSICIAN IMMEDIATELY.

UNDESIRABLE AND UNPLEASANT EFFECTS

As with any medicine, this product may produce unpleasant effects of varying severity in some people.

Tell your physician about any unwanted or unpleasant effect which is not mentioned in this information leaflet.

INSTRUCTIONS

Store below 25°C. Protect from heat and light. Keep all medicines out of the reach of children. To be sold on the prescription of a registered medical practitioner only.

PRESENTATION

Tri-D Injection 5 x 1ml ampoule.

ٹرائی-ڈی انجکشن
(کول کلسی فرول)

ہدایات: دوا کو 25° سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔ گرمی اور روشنی سے محفوظ رکھیں۔
تمام دوائیں بچوں کی پہنچ سے دُور رکھیں۔



Manufactured by:
STANDPHARM PAKISTAN (PVT) LTD
20 km Ferozepur Road Lahore, Pakistan.